

PARAMOL®
120mg/ 5mL Syrup

Dear patient,

Please read the following instructions carefully. They contain important information about the use of this medicine. If you have any further questions, please ask your doctor or pharmacist.

Information about PARAMOL

Each 5 mL of PARAMOL syrup contains 120 mg paracetamol.

The excipients are: Citric acid, sodium benzoate, glycerin, sucrose, sodium edetate, red erythrosine, strawberry, menthol and purified water.

Paracetamol is a safe and effective analgesic and antipyretic drug that is used to temporarily relieve pain and fever.

PARAMOL syrup is indicated for babies and infants in the following conditions:

- Sore throat
- Toothache
- Teething
- Childhood infections
- Immunization

The way to take PARAMOL

PARAMOL syrup must be administered as directed by the doctor. The dose is generally individualized according to the child's weight or age.

A dose of 10-15 mg/kg to be repeated every 4-6 hours if necessary.

Weight (kg)	Age	Dose (mL)
5 – 6	2-3 months	3
6 – 7	3-4 months	3.5
7 – 8	4-6 months	4
8 – 9	6-8 months	5
9 - 10	8-12 months	5.5
10 – 12	1-2 years	6
12 - 14	2-3 years	7.5
14 - 16	3-4 years	8.5
16 - 18	4-5 years	10

- The dose may be given directly in the mouth or mixed with a little juice or water first.

In case of overdose

In case of intake of high doses of this medication, inform the doctor at once and seek emergency medical attention. General measures should be adopted.

In case of missed dose

Give the missed dose as soon as you remember unless the next intake is near. Go on giving the next scheduled dose as directed. Do not take a double dose at once.

Contraindications

This drug is contraindicated in case of hypersensitivity to Paracetamol or to any of the components.

Precautions

- Inform your doctor before giving PARAMOL in case your child has glutathione depleted states such as sepsis, or kidney or liver problems
- Do not give to your child more than 60 mg /kg presented in divided doses throughout 24 hours
- Do not give to your child for longer than 3 days without asking a doctor
- Contact your doctor if fever persists after the second dose
- For other indications, give only under medical advice. It is not recommended for children under 2 months

Associations with other medications

Please inform your doctor or pharmacist if other medicines are being taken or have been taken recently.
Do not give PARAMOL with other products containing paracetamol.

Adverse reactions

This drug is generally well tolerated when used as directed.

Storage

Store at controlled room temperature (up to 30°C), protected from light and humidity, beyond the reach of children.

The expiry date is printed on the pack; don't use this medicine after this date.

Pack Presentation

PARAMOL, paracetamol 120mg / 5mL, bottle of 60 mL with a dosing cup

Revision date: 03/2023

PARS1/003