

HISTAMED-F® Syrup

Dear patient,

Please read the following instructions carefully. They contain important information about the use of this medicine. If you have any further questions, please ask your doctor or pharmacist.

Information about HISTAMED-F Syrup

Each 5 mL of HISTAMED-F syrup contains:

Dexamethasone 0.25 mg

Dexchlorpheniramine maleate 2 mg

HISTAMED-F syrup is for oral use.

Dexamethasone is a synthetic glucocorticoid having potent anti-inflammatory effects in disorders of many organ systems.

Dexchlorpheniramine is an H1-receptor antagonist. It is an alkylamine derivative. Dexchlorpheniramine antagonizes most of the pharmacological effects of histamine.

HISTAMED-F is effective for the prevention and relief of many allergic manifestations.

HISTAMED-F is indicated for the treatment of allergic and inflammatory conditions:

- Perennial and seasonal allergic rhinitis: HISTAMED-F relieves sneezing, rhinorrhea, nasal itch and conjunctivitis
- Acute and chronic allergic and inflammatory processes involving the eye such as allergic conjunctivitis, keratitis, nongranulomatous iritis, iridocyclitis, chorioretinitis and uveitis.
- Allergic skin disorders including mild, uncomplicated, acute urticaria and angioedema, pruritus, insect bites, some drug allergies, atopic dermatitis (eczema), and contact dermatitis
- Allergic and non-allergic pruritic symptoms
- Adjunctive therapy in anaphylactic reactions
- Severe chronic asthma and seasonal asthma

The way to take HISTAMED-F SYRUP

Take HISTAMED-F Syrup as directed.

Do not discontinue the treatment without consulting your doctor.

Dosage is individualized on the basis of the condition under treatment and the response of the patient.

Unless otherwise directed by your doctor, the recommended average initial doses are:

Children 3 to 6 years of age	half a teaspoonful (2.5 ml) or 1 teaspoonful (5 ml) once daily
Children 6 to 12 years of age	1 teaspoonful (5 ml) once or twice daily
Adults and children over 12 years of age	1 teaspoonful (5 ml) 4 times daily, after meals and at bedtime

In severe cases this dosage can be increased up to 6 teaspoonfuls of the syrup a day. However, as improvement takes place the dosage should be reduced to the minimum maintenance level.

In younger children, the dosage should be adjusted according to the severity of the condition treated rather than by age or weight.

Duration

The duration of the treatment is determined according to the condition under treatment.

In ordinary self-limited conditions, e.g. poison ivy dermatitis, simple urticaria and certain cases of contact dermatitis, a short course of therapy will relieve symptoms and may in these cases be considered as a cure course.

In seasonal allergic disorders, such as hay fever, asthma and allergic conjunctivitis, therapy will not be required during times when the offending allergen is not present in the patient's environment.

Certain chronic disorders resistant to standard allergy treatment may require prolonged maintenance therapy (eczema, perennial rhinitis, chronic urticaria).

In case of overdose

In case of intake of high doses of this medication, inform your doctor at once and seek emergency medical attention. General measures should be adopted.

In case of missed dose

Take the missed dose as soon as you remember unless the next intake is near. Go on taking the next scheduled dose as directed. Do not take a double dose at once.

Contraindications

This drug is contraindicated in the following cases:

- history of hypersensitivity to any of the components
- greatly depressed states due to central nervous system depressants, bone marrow depression
- viral infections

Precautions

- In case of pregnancy, consult your doctor about the use of this medicine. This drug should not be used during lactation.
- Caution must be used in case of liver or renal diseases.
- This drug may cause marked drowsiness or impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a vehicle or operating machinery.
- This drug must be used with caution in elderly patients and in patients with a predisposition to urinary retention, increased intraocular pressure, hyperthyroidism, diabetes, chronic constipation, cardiovascular disease, hypertension, osteoporosis, tuberculosis or gastroduodenal ulcer.

Associations with other medications

Please inform your doctor if other medicines are being taken or have been taken recently.

Use with caution with alcohol, hypnotics, narcotic analgesics, oral anticoagulants, antidepressants, tranquilizers, anti-arrhythmic agents, atropine-like substances, digitalis glycosides, carbamazepine, phenobarbital, phenytoin, primidone, rifampicine, antidiabetics, isoniazide and bromocriptine.

Adverse reactions

This drug is usually well tolerated when used as directed.

Side effects related to the presence of the corticoid may appear during a prolonged use or at high doses.

Certain patients may, however, respond to antihistamine drugs by becoming drowsy, dizzy or nauseated. Other possible side effects include restlessness, fatigue, dry mouth, anorexia, blurred vision, constipation and headache.

Inform your doctor if any side effect appears or becomes bothersome.

Storage

Store at controlled room temperature (up to 30°C), protected from light and humidity, beyond the reach of children.

The expiry date is printed on the pack; don't use this medicine after this date.

Pack Presentation

HISTAMED-F syrup, bottle of 90 mL with a dosing cup.

Revision date: 01/2009

HISS/003